

## **Controlling Purple Jewelweed (*Impatiens glandulifera*): Assessment of Feasibility and Costs**

Author(s): Michaël Leblanc and Claude Lavoie

Source: Invasive Plant Science and Management, 10(3):254-261.

Published By: Weed Science Society of America

URL: <http://www.bioone.org/doi/full/10.1017/inp.2017.21>

---

BioOne ([www.bioone.org](http://www.bioone.org)) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/page/terms\\_of\\_use](http://www.bioone.org/page/terms_of_use).

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

# Controlling Purple Jewelweed (*Impatiens glandulifera*): Assessment of Feasibility and Costs

Michaël Leblanc and Claude Lavoie\*

We experimentally tested the feasibility of a control campaign of purple jewelweed (*Impatiens glandulifera*), an exotic invasive species in Europe and North America. We evaluated the amount of time and money required to control the plant along riverbanks, with particular attention paid to the recovery of riparian vegetation following hand pulling and bagging. Work time was directly and significantly related to stem density and fresh biomass of the invader, but the relationship was stronger for density. Density and biomass were strongly reduced by the first hand-pulling operation from a mean of 45 to 2 stems m<sup>-2</sup> and from a mean of 0.95 kg m<sup>-2</sup> to nearly zero, a good performance but not enough to negate the need for a second hand pulling later in the summer. A single hand pulling significantly reduced the cover of purple jewelweed from 30% to 7%. Riparian vegetation disturbed by the first hand pulling largely recovered during the following 30 d. Expressed over an area of 1 ha, the total amount of time required to control purple jewelweed is 1,400 work hours over 2 yr, or a minimum investment of Can\$21,000 (US\$17,000). Although controlling a well-established purple jewelweed population is expensive, to properly evaluate the benefits, we must also consider the costs of soil erosion caused by this species.

**Nomenclature:** Policeman's helmet, purple jewelweed, *Impatiens glandulifera* Royle.

**Key words:** Control campaign cost, erosion, hand pulling, riparian vegetation, river.

Substantial amounts of scientific information on invasive species are available to environmental managers. Unfortunately, managers rarely rely on the scientific literature for developing weed control programs, largely because there is a mismatch between the type of research conducted by invasion biologists (essentially fundamental) and the needs of practitioners (essentially applied), a clear example of the so-called “knowing–doing gap” (Lavoie and Brisson 2015; Matzek et al. 2014, 2015). Reliable information on the cost of control campaigns is especially lacking, which complicates the task of managers who must evaluate the costs and benefits of different control methods and the feasibility of large-scale operations conducted with limited resources (Braun et al. 2016; Delbart et al. 2012).

Purple jewelweed (Balsaminaceae: *Impatiens glandulifera*; hereafter PUJE) is one of the invasive species for which a significant amount of information is available on the biology and ecology of the plant, but little information is available

on the control methods and their associated costs. The native range of this annual vascular plant is located in the western part of the Himalayas (India, Pakistan), where it can grow up to an altitude of 4,000 m (Clements et al. 2008). An individual may reach a height of 2.5 m at midsummer. Although its root system is shallow, the large stem base and the adventitious roots help solidly anchor the plant in the soil (Ennos et al. 1993). A single plant may produce 800 to 2,500 seeds contained in capsules that split explosively at maturity, dispersing the seeds up to 5 to 6 m. (Beerling and Perrins 1993; Chapman and Gray 2012; Clements et al. 2008). Beerling and Perrins (1993) argued that PUJE seeds can, under certain circumstances, remain viable in the soil for at least 18 mo, but Perglová et al. (2009) concluded, on the basis of laboratory and garden experiments, that a seedbank is unlikely for this species. Seeds can be dispersed by water, especially during floods (Čuda et al. 2017; Love et al. 2013; Pyšek and Prach 1995), but the range of distances achieved by water dispersal along streams and rivers is unknown.

PUJE was introduced in Europe in 1839 and in north-eastern North America in 1883 (Perrins et al. 1993; Tabak and von Wettberg 2008). It has spread rapidly along rivers and drainage ditches, where it can form dense stands

DOI: 10.1017/inp.2017.21

\* First and second authors: Graduate Student and Professor, École Supérieure d'Aménagement du Territoire et de Développement Régional, Université Laval, Québec City, QC G1A 0V6, Canada. Corresponding author's E-mail: [claudio.lavoie@esad.ulaval.ca](mailto:claudio.lavoie@esad.ulaval.ca)

## Management Implications

Controlling a well-established population of purple jewelweed (*Impatiens glandulifera*) on riverbanks is probably unrealistic on a regional scale, but a local, systematic (from upstream to downstream), hand-pulling control is technically practicable along a river system. To locally extirpate purple jewelweed, two hand-pulling operations per summer repeated over 2 yr are effective, so long as the controlled area is double-checked during each operation for missed plants, especially flowering individuals. The riparian vegetation may spontaneously recover following an early summer hand-pulling campaign, particularly if grass species with rhizomatous root systems are present. However, controlling purple jewelweed is expensive: a 1-km stretch of a brook (two riverbanks 5- to 6-m wide), requires 1,500 to 1,600 work hours, a minimum investment of Can\$24,000 (US\$18,000). Targeting small populations before they spread into available habitats would be more economical. On the other hand, dredging a 1-km section of a brook to remove an accumulation of sediments may represent costs of about Can\$30,000 (US\$23,000). Considering purple jewelweed may accelerate riverbank erosion, what initially appears to be a costly control campaign could in fact represent a wise investment to prevent recurrent problems.

(Dawson and Holland 1999; Kostrakiewicz-Gierałt and Zając 2014; Perrins et al. 1993; Pyšek and Prach 1995). The effects of PUJE on the biodiversity of riparian habitats is nevertheless controversial. Several studies have shown that the invader has a low impact on plant species richness and composition (Diekmann et al. 2016; Hejda and Pyšek 2006; Hejda et al. 2009) and that the invaded habitats even support a higher richness and abundance of gastropods (Ruckli et al. 2013). Others showed that PUJE may change the composition of snail and microarthropod assemblages (Horáčková et al. 2014; Rusterholz et al. 2014) or reduce the abundance of foliage- and ground-dwelling invertebrates (Tanner et al. 2013). Some ecologists found evidence that PUJE outcompetes native plants for pollinators and disrupts pollen transfer networks (Chittka and Schürkens 2001; Lopezaraiza-Mikel et al. 2007; Thijs et al. 2012), but others concluded that there was no impact on pollination (Bartomeus et al. 2010; Emer et al. 2015; Vilà et al. 2009). A serious concern is soil erosion, a consequence of the replacement of perennials by PUJE, which leaves riverbanks devoid of vegetation cover after plant death and decomposition in late fall. Soil losses may reach, in heavily invaded riparian sites, 2,300,000 kg (Greenwood and Kuhn 2014). Greenwood and Kuhn (2015) estimated that the increasing dominance of PUJE in riverbanks could compromise their ability to provide certain ecosystem services, such as filtering nutrient-rich sediments, moderating stream flow, and providing natural flood protection.

The scientific literature on the management and control of PUJE is sparse. Models predicted that removing upstream populations first, in an effort to locally reduce the ability of

the species to colonize downstream parts of river systems, could be effective. However, from a regional point of view, a 99% control efficiency over a 25-yr period would be almost as ineffective as no management at all with regard to controlling the species (Wadsworth et al. 2000). At the Thayatal-Podyjí National Park (Austria–Czech Republic), repeated mowing and hand-pulling operations allowed a major (87%) reduction of the area covered by PUJE populations within a decade (Schiffleithner and Essl 2016). To our knowledge, the only experiment comparing the respective efficacy of PUJE control techniques (Cockel et al. 2014) showed that hand pulling was much more effective than mowing for reducing the PUJE cover over a 2-yr period. In summary, the control of a well-established population of PUJE is probably unrealistic from a regional point of view, but the local, systematic (from upstream to downstream), hand pulling of PUJE is apparently technically practicable along a river system. However, the question remains: Is such an operation cost-effective?

We tested the feasibility of a PUJE control (hand-pulling) campaign conducted along a small brook strongly invaded by the species. We evaluated the amount of time required to control the plant, i.e., to reduce the number of individuals to the lowest possible level, as close as possible to complete extirpation, according to density and biomass. We paid particular attention to the disposal of pulled plants, which must be considered in the total cost, and to the rapid recovery of riparian vegetation following hand pulling, a critical point for the prevention of soil erosion and the opportunistic establishment of other invasive species that could negate the benefits of PUJE removal (Cockel et al. 2014; Hulme and Bremner 2006). We accurately calculated the cost of a control campaign. We hypothesized that PUJE removal time would be directly proportional to stem density and biomass, and that a hand-pulling campaign beginning in early summer would allow a spontaneous (unassisted) and complete recovery of the riparian vegetation. However, we also predicted that a PUJE control campaign would be extremely expensive where the species is already well established.

## Materials and Methods

**Study Area.** The PUJE population selected for this study was located along the Sainte-Geneviève Brook (hereafter SGBk), at Saint-Isidore, near Québec City, in Québec, Canada (46.98°N, 71.15°W). SGBk is the main tributary of the Fourchette Brook, which flows into the Etchemin River that ends its course in the St Lawrence River at the town of Lévis, across the river from Québec City. The mean annual temperature in the study area is 5 C, the mean temperature of the coldest month (January) is -12 C, and that of the warmest month (July) is 19 C. The mean annual precipitation totals 1,178 mm, 23% of which falls as snow (Government of Canada 2017).



Figure 1. Purple jewelweed (*Impatiens glandulifera*) population at Sainte-Geneviève Brook (Québec, Canada). Photograph taken August 16, 2016 (M. Leblanc).

SGBk is 11-km long. The brook, 2- to 3-m wide, is bordered by agricultural lands (corn, soybean) over 80% of its course. Riverbanks have an average inclination of 36°. They are essentially clayey or loamy, with some rocky sections, and are covered by a herbaceous riparian strip about 5- to 6-m wide in cultivated areas. Province of Québec legislation requires the preservation of a vegetated riparian strip at least 3-m wide on both sides of a stream in agricultural areas (Loi sur la qualité de l'environnement, chapitre Q-2, r. 35). At the study site, the riverbank vegetation is largely dominated by PUJE and two other perennial exotic invasive species, reed canarygrass (*Phalaris arundinacea* L.) and smooth brome (*Bromus inermis* Leyss.), with almost no shrubs nor trees.

About 2001 (exact introduction year unknown), some PUJE individuals were planted a few meters from SGBk, in a residential garden located 7.4 km from the source of the brook. PUJE has since spread along SGBk: in 2016 the riverbanks were heavily infested (up to 177 individuals  $m^{-2}$ ) downstream from the introduction point (Figure 1). Several thousands of individuals were also detected along the course of the Fourchette Brook, from its junction with SGBk to the Etchemin River (C Lavoie, unpublished data).

**Data Sampling.** We selected a 1-km-long stretch of SGBk, 535 m downstream from the introduction point of the species, at the beginning of a cultivated zone where conditions were uniform with regard to width and composition of the vegetated riparian strip and adjacent land use. A hand-pulling station was systematically located every 50 m along the stretch, on each riverbank, for a total 42 stations; three of these were subsequently rejected because they were too heavily disturbed by recent agricultural activities

(vegetation in large part removed). Before hand pulling began, a 24- $m^2$  area was delimited at each station, 4-m wide by 6-m long (32 stations), or 3-m wide by 8-m long where the riparian strip was narrower (7 stations). The following data were taken: (1) the width of the riparian strip; (2) the width and (3) the water level of the brook; and (4) the slope of the riverbank. The vegetation cover inside the 24- $m^2$  area was sampled just before the first and the second hand pulling using a point intercept method (Meunier and Lavoie 2012). Lines were drawn 1-m apart across each station. The presence of four plant categories touching a 3-cm-wide rod (held perpendicular to ground) was recorded at every meter along each line: (1) PUJE; (2) grass species; (3) forb species others than PUJE; and (4) woody species. The presence of the other invasive species (reed canarygrass and smooth brome) was also noted. This method provided frequency data (indicator of plant cover) from 35 or 36 sampling points per station. Photographs were taken just before and after the first hand pulling and before the second hand pulling (Figure 2). The total experimental area in 2016 was 936  $m^2$ .

As previously mentioned, two hand-pulling operations were conducted in summer 2016, the first from June 30 to July 14, when most PUJE individuals reached a height of about 60 cm and were thus easy to detect and hand pull, and the second from August 15 to 18, to detect individuals that were not eliminated during the first hand pulling and document the recovery of the vegetation of the riparian strip. Hand pulling was conducted by two or three wildlife technicians (student trainees). Pulled plants were dumped in 140-L trays that were emptied as needed on a tarp. During the first hand pulling, technicians made a single attempt to remove an individual with its root system; if the stem broke without roots, the base of the stem was left on site. This procedure was used to evaluate whether a quicker and cheaper hand-pulling operation would be as effective as a meticulous (all roots removed) but more time-consuming protocol. However, during the second hand pulling, all stems were removed with their root systems, even if the stems initially broke off. After the end of each hand pulling, the area was double-checked for missed stems. Extracted plants were tallied by presence or absence of root system and flowers. Fresh biomass was determined with a portable scale. For this experiment, fresh biomass was more relevant than dry biomass, given that technicians were handling fresh tissue. Pulled plants were packed into black garbage bags to prevent rerooting, and left on site to die by solarization. Time required for hand pulling and bagging was noted.

**Data Analyses.** The relationship between PUJE stem density or biomass and hand-pulling time was analyzed with linear regressions. Wilcoxon matched-pairs signed-rank tests were used to detect significant differences between the vegetation cover before the first and the second hand




Figure 2. Overview of two sampling stations on the riverbanks of Sainte-Geneviève Brook (Québec, Canada) just before (A: July 5, 2016, 177 stems  $m^{-2}$ ; B: July 14, 2016, 83 stems  $m^{-2}$ ) the first hand pulling of purple jewelweed (*Impatiens glandulifera*), just after (C, D) the first hand pulling, and just before (E: August 16, 2016, 3 stems  $m^{-2}$ ; F: August 18, 2016, 3 stems  $m^{-2}$ ) the second hand pulling. (Photographs: M Leblanc.).

pulling, considering data were not normally distributed (Scherrer 1984). Stata software (StataCorp 2013) was used for calculations.

Operation costs were estimated for a 1-km stretch of a brook and over a 2-yr period. For Year 1, we calculated the total time (two hand-pulling operations) required to extract and pack PUJE stems for the area treated in 2016 (936  $m^2$ ). We then extrapolated this time to the total riparian strip area of SGBk over 1 km, estimated using the mean value of the width of the strips measured at the 39 hand-pulling stations.

On the basis of previous experiments (Cockel et al. 2014; Schiffeithner and Essl 2016) and the results of this study,

we concluded that a single summer of hand pulling would not be sufficient for an effective control of PUJE. Some small stems, hidden in the vegetation, will inevitably be missed by technicians in Year 1. Although most are unlikely to reach maturity by the end of the summer, the few that do will produce seeds. Our experimental design did not allow a second year of hand pulling in 2017, because the stations were reseeded by the thousands of bordering PUJE individuals; in a real control procedure, all individuals would be systematically eliminated upstream to downstream to prevent reseeding. So we estimated that it would be necessary to conduct a second summer of control (Year 2),

with one hand pulling at the beginning of the summer and an additional one at midsummer. We estimated that the effort of each hand pulling at Year 2 would be equivalent to the time required to conduct the second hand pulling of 2016 (Year 1). In fact, technicians will probably spend more time searching for residual PUJE stems than actually hand pulling in Year 2.

The cost of the labor force was calculated using the mean hourly wage (including social benefits) of a wildlife technician hired in Québec, which was Can\$33.39 in 2015, i.e., about US\$26 (Institut de la statistique du Québec 2016), or the hourly wage (with benefits) paid to the students who were hired for this study (Can\$15.16, US\$12), which would represent the minimum wage that can reasonably be paid under these circumstances.

## Results and Discussion

The density and biomass of PUJE stems before the first hand pulling was highly variable across the 39 stations investigated along SGBk (Table 1), from 1 to 177 stems  $m^{-2}$  (mean: 45 stems  $m^{-2}$ ) and from a value close to zero to 3.55 kg  $m^{-2}$  (mean: 0.95 kg  $m^{-2}$ ). Stem density (mean value) was higher at SGBk than in the native range of the species (India: 18 to 30 stems  $m^{-2}$ ; Tanner et al. 2014) but very close to values recorded in Europe (Beerling and Perrins 1993; Greenwood and Kuhn 2014). The mean frequency of PUJE (indicator of plant cover) was 30% before the first hand pulling (Figure 3), varying from 3% to 77% among stations, which was roughly the same as recorded in various riparian habitats of the Czech Republic and Germany (Bartomeus et al. 2010; Hejda and Pyšek 2006). Higher covers (60% to 100%) have nevertheless been observed in Europe (Cockel et al. 2014; Emer et al. 2015; Hejda et al. 2009; Hulme and Bremner 2006). Although invaded by PUJE, the SGBk riverbanks maintained a high cover of grass species (mean frequency: 87%), of which nearly half (43%) was smooth brome.

Hand-pulling and bagging time was directly and significantly related to stem density and biomass, but the relationship was stronger for density (Figure 4). Evaluating stem density prior to the first hand pulling with the

following equation can thus provide a good estimation of the time required for managing the operation:

$$t = (1.7d + 54)/24 \quad [1]$$

where  $t$  is the total amount of time (in minutes) required for extirpating PUJE over an area of  $1 m^2$ , and  $d$  is the PUJE stem density per square meter.

Only 10% of the stems were pulled without their root system during the first hand pulling, a good performance but not enough to negate the need for a second hand pulling during the same summer. Density and biomass were strongly reduced by the first hand pulling to a mean of 2 stems  $m^{-2}$  and to a value close to zero for biomass (Table 1), but it should be noted that a certain number of stems would also have been naturally thinned through competition for light and space, especially in the stations with the highest density (Beerling and Perrins 1993). About 14% of the stems pulled at mid-August were flowering and could have produced a new generation of seeds if not removed. However, it is likely this proportion would have been lower if all stems detected during the first hand pulling had been removed with their root systems (Clements et al. 2008).

The first hand pulling significantly ( $P < 0.0001$ ) reduced the cover (frequency) of PUJE from 30% to 7% (mean values; Figure 3) and had a significant effect on the forb (reduction;  $P = 0.0211$ ) and grass (increase;  $P = 0.0068$ ) cover, although changes were small. In fact, the plant cover disturbed by the first hand pulling largely recovered during the following 30 d (Figure 2), because of the vegetative spread (rhizomes) of grass species, particularly reed canary-grass and smooth brome. Reed canarygrass and smooth brome are highly invasive species (Lavoie et al. 2005; Otfinowski et al. 2007; Stannard and Crowder 2003), and this probably explains the rapid recovery of the riparian vegetation, although other experiments also showed that riparian vegetation may rapidly recover within 1 mo of PUJE removal, because most riverbank species have a seedbank or vegetative dispersal means (Hulme and Bremner 2006).

All stations considered, the first hand pulling took six times longer than the second (Table 1). A total of 101 work hours was required to remove PUJE stems from 936  $m^2$ ; 9%

Table 1. Working time required in summer 2016 to hand pull and bag purple jewelweed from 39 sampling stations covering a total area of 936  $m^2$  on the riverbanks of Sainte-Geneviève Brook (Québec, Canada).

Hand pulling	Total time in work hours	Stem density — $n m^{-2}$ —			Fresh biomass —kg $m^{-2}$ —		
		Mean $\pm$ SD	Minimum	Maximum	Mean $\pm$ SD	Minimum	Maximum
1 (June 30–July 14)	86	45 $\pm$ 46	1	177	0.95 $\pm$ 1.03	0.02	3.55
2 (August 15–18)	15	2 $\pm$ 3	0	15	0.03 $\pm$ 0.04	0.01	0.17




Figure 3. Frequency data (indicator of plant cover; mean  $\pm$  SD) of four plant categories in sampling stations on the riverbanks of Sainte-Geneviève Brook (Québec, Canada), just before the first hand pulling of purple jewelweed (*Impatiens glandulifera*) conducted from June 30 to July 14, 2016, and just before the second hand pulling conducted from August 15 to 18, 2016.

of this time was used for bagging. For a 1-km stretch of a brook (two riverbanks), under SGBk conditions (mean bank width: 5.6 m), this represents 1,208 h for the first year (Table 2). But since a second year would probably be necessary to extirpate the plant, the total amount of time required to control PUJE over this distance totals 1,566 h. This represents 1,398 work hours  $\text{ha}^{-1}$ . By comparison, 270 work hours  $\text{ha}^{-1}$  over a 10-yr period were necessary to reduce PUJE extent and density by 77% at the Thayatal-Podyjí National Park (Schiffleithner and Essl 2016). The context and methods were different, and the PUJE cover was much lower (1% to 3%) in the park than at SGBk (F Essl, personal communication), which is a precaution against comparing worker efficiency, but both cases provide insight on the magnitude of effort required to control PUJE stands.

In the Québec context, controlling (hand pulling and bagging) PUJE from SGBk over a 1-km distance would represent a minimum investment of Can\$23,742 (about US\$18,000), but could reach Can\$52,289 (about US \$40,000) if experienced (higher wages) wildlife technicians were employed (Table 2). This is probably a minimum estimation, since we cannot exclude the possibility that




Figure 4. Duration of purple jewelweed (*Impatiens glandulifera*) hand-pulling operations, including bagging (1: from June 30 to July 14, 2016; 2: from August 15 to 18, 2016), according to stem density and fresh biomass, conducted in 39 sampling stations (area: 24 m<sup>2</sup>) on the riverbanks of Sainte-Geneviève Brook (Québec, Canada).

Table 2. Estimation of the cost of a control campaign of purple jewelweed along 1 km of riverbanks (1,1 ha) and over 2 yr in Québec (Canada).

Year	Hand pulling	Hand pulling —work hours—	Bagging	Total	Labor force cost: wildlife technician (hand pulling + bagging)			
					Minimum wage <sup>a</sup>		Mean wage <sup>b</sup>	
					Can\$	US\$	Can\$	US\$
1	1	921	108	1029	15,600	11,932	34,358	26,280
	2	177	2	179	2714	2076	5977	4572
2	1	177	2	179	2714	2076	5977	4572
	2	177	2	179	2714	2076	5977	4572
Total		1452	114	1566	23,742	18,160	52,289	39,996

<sup>a</sup> Can\$15.16 (US\$12), social benefits included: wage paid to student technician trainees in 2016.

<sup>b</sup> Can\$33.39 (US\$26), social benefits included (according to Institut de la statistique du Québec 2016).

more than 2 yr may be necessary to extirpate the plant. Bagging was efficient—stems rapidly decomposed in the bags—but to what extent this procedure was really essential to prevent rerooting remains unclear. However, not bagging would only reduce costs by 7%. We contacted 25 non-governmental organizations in Québec who are in charge of the management and environmental protection of watersheds and the control of riparian invaders, and none (with one exception) would be able to allocate such a sum for controlling PUJE without municipal or state (provincial) support, which is rarely provided in Québec for controlling invasive species.

Controlling a well-established (about 15-yr-old) PUJE population is very expensive, and as highlighted by Meier et al. (2014), targeting small stands before they spread into available habitats would be more economical. Other control options (burning, herbicides, tilling) are strictly prohibited in Québec in riparian habitats (Loi sur la qualité de l'environnement, chapitre Q-2, r. 35; Loi sur les pesticides, chapitre P-9.3, r. 1) and cannot be considered to reduce labor costs. On the other hand, the same stretch of SGBk we used in our experiments had to be dredged in 2015 to remove an accumulation of sediments impeding drainage and causing flooding in the Saint-Isidore village. This cost Can\$30,000 (about US\$23,000) per kilometer. Without additional supporting data, we cannot prove this was the result of accelerated riverbank erosion due to an extensive PUJE infestation, but what initially appears to be a costly control campaign could in fact represent a wise investment to prevent recurrent problems.

### Acknowledgments

This research was financially supported by the Natural Sciences and Engineering Research Council of Canada. We are grateful to the Conseil de bassin de la rivière Etchemin for

having provided field work assistance (Clovis Arguin and Jacob Carrier), to Elisabeth Groeneveld and Nicolas Trottier for additional help on this project, to Steeven Lagrange (Steegrain, Inc.) and Christian Larochelle (Laroporc, Inc.) for access to the study site, and to two anonymous reviewers for comments on an earlier draft.

### Literature Cited

- Bartomeus I, Vilà M, Steffan-Dewenter I (2010) Combined effects of *Impatiens glandulifera* invasion and landscape structure on native plant pollination. *J Ecol* 98:440–450
- Beerling DJ, Perrins JM (1993) *Impatiens glandulifera* Royle (*Impatiens roylei* Walp.). *J Ecol* 81:367–382
- Braun M, Schindler S, Essl F (2016) Distribution and management of invasive alien plant species in protected areas in Central Europe. *J Nat Conserv* 33:48–57
- Chapman DS, Gray A (2012) Complex interactions between the wind and ballistic seed dispersal in *Impatiens glandulifera* (Royle). *J Ecol* 100:874–883
- Chittka L, Schürkens S (2001) Successful invasion of a floral market. An exotic Asian plant has moved in on Europe's river-banks by bribing pollinators. *Nature* 411:653
- Clements DR, Feenstra KR, Jones K, Staniforth R (2008) The biology of invasive alien plants in Canada. 9. *Impatiens glandulifera* Royle. *Can J Plant Sci* 88:403–417
- Cockel CP, Gurnell AM, Gurnell J (2014) Consequences of the physical management of an invasive alien plant for riparian plant species richness and diversity. *River Res Appl* 30:217–229
- Čuda H, Rumlerová Z, Brůna J, Skálová H, Pyšek P (2017) Floods affect the abundance of invasive *Impatiens glandulifera* and its spread from river corridors. *Divers Distrib* 23:342–354
- Dawson FH, Holland D (1999) The distribution in bankside habitats of three alien invasive plants in the U.K. in relation to the development of control strategies. *Hydrobiologia* 415:193–201
- Delbart E, Mahy G, Weickmans B, Henriët F, Crémer S, Pieret N, Vanderhoeven S, Monty A (2012) Can land managers control Japanese knotweed? Lessons from control tests in Belgium. *Environ Manag* 50:1089–1097



- Diekmann M, Effertz H, Baranowski M, Dupré C (2016) Weak effects on plant diversity of two invasive *Impatiens* species. *Plant Ecol* 217:1503–1514
- Emer C, Vaughan IP, Hiscock S, Memmott J (2015) The impact of the invasive alien plant, *Impatiens glandulifera*, on pollen transfer networks. *PLoS ONE* 10:e0143532
- Ennos AR, Crook MJ, Grimshaw C (1993) A comparative study of the anchorage systems of Himalayan balsam *Impatiens glandulifera* and mature sunflower *Helianthus annuus*. *J Exp Bot* 44:133–146
- Government of Canada (2017) Historical Climate Data. [http://climate.weather.gc.ca/index\\_e.html](http://climate.weather.gc.ca/index_e.html). Accessed: January 1, 2017
- Greenwood P, Kuhn NJ (2014) Does the invasive plant, *Impatiens glandulifera*, promote soil erosion along the riparian zone? An investigation on a small watercourse in northwest Switzerland. *J Soil Sediment* 14:637–650
- Greenwood P, Kuhn NJ (2015) Commentary on “Plants as river system engineers” by A. Gurnell. *Earth Surface Processes and Landforms* 39:4–25, 2014. DOI 10.1002/esp.3397. *Earth Surf Proc Land* 40:131–134
- Hejda M, Pyšek P (2006) What is the impact of *Impatiens glandulifera* on species diversity of invaded riparian vegetation? *Biol Conserv* 132:143–152
- Hejda M, Pyšek P, Jarošík V (2009) Impact of invasive plants on the species richness, diversity and composition of invaded communities. *J Ecol* 97:393–403
- Horáčková J, Juříčková L, Šizling AL, Jarošík V, Pyšek P (2014) Invasiveness does not predict impact: response of native land snail communities to plant invasions in riparian habitats. *PLoS ONE* 9:e108296
- Hulme PE, Bremner ET (2006) Assessing the impact of *Impatiens glandulifera* on riparian habitats: partitioning diversity components following species removal. *J Appl Ecol* 43:43–50
- Institut de la statistique du Québec (2016) Résultats de l'Enquête sur la rémunération globale au Québec. Collecte 2015. <http://www.stat.gouv.qc.ca/statistiques/travail-remuneration/resultats-erg-2015.pdf>. Accessed: February 13, 2017
- Kostrakiewicz-Gierałt K, Zajac M (2014) The influence of habitat conditions on the performance of two invasive, annuals: *Impatiens glandulifera* and *Bidens frondosa*. *Biologia* 69:449–462
- Lavoie C, Brisson J (2015) Training environmental managers to control invasive plants: acting to close the knowing-doing gap. *Invasive Plant Sci Manag* 8:430–435
- Lavoie C, Dufresne C, Delisle F (2005) The spread of reed canarygrass (*Phalaris arundinacea*) in Québec: a spatio-temporal perspective. *Écoscience* 12:366–375
- Lopezaraiza-Mikel ME, Hayes RB, Whalley MR, Memmott J (2007) The impact of an alien plant on a native plant–pollinator network: an experimental approach. *Ecol Lett* 10:539–550
- Love HM, Maggs CA, Murray TE, Provan J (2013) Genetic evidence for predominantly hydrochoric gene flow in the invasive riparian plant *Impatiens glandulifera* (Himalayan balsam). *Ann Bot* 112:1743–1750
- Matzek V, Covino J, Funk JL, Saunders M (2014) Closing the knowing-doing gap in invasive plant management: accessibility and inter-disciplinarity of scientific research. *Conserv Lett* 7:208–215
- Matzek V, Pujaleat M, Cresci S (2015) What managers want from invasive species research versus what they get. *Conserv Lett* 8:33–40
- Meier ES, Dullinger S, Zimmermann NE, Baumgartner D, Gattringer A, Hülber Karl (2014) Space matters when defining effective management for invasive plants. *Divers Distrib* 20:1029–1043
- Meunier G, Lavoie C (2012) Roads as corridors for invasive plant species: new evidence from smooth bedstraw (*Galium mollugo*). *Invasive Plant Sci Manag* 5:92–100
- Ostrowski R, Kenkel NC, Catling PM (2007) The biology of Canadian weeds. 134. *Bromus inermis* Leyss. *Can J Plant Sci* 87:183–198
- Perglová I, Pergl J, Skálová H, Moravcová L, Jarošík V, Pyšek P (2009) Differences in germination and seedling establishment of alien and native *Impatiens* species. *Preslia* 81:357–375
- Perrins J, Fitter A, Williamson M (1993) Population biology and rates of invasion of three introduced *Impatiens* species in the British Isles. *J Biogeogr* 20:33–44
- Pyšek P, Prach K (1995) Invasion dynamics of *Impatiens glandulifera*: a century of spreading reconstructed. *Biol Conserv* 74:41–48
- Ruckli R, Rusterholz H-P, Baur B (2013) Invasion of *Impatiens glandulifera* affects terrestrial gastropods by altering microclimate. *Acta Oecol* 47:16–23
- Rusterholz H-P, Salamon J-A, Ruckli R, Baur B (2014) Effects of the annual invasive plant *Impatiens glandulifera* on the Collembola and Acari communities in a deciduous forest. *Pedobiologia* 57:285–291
- Scherrer B (1984) *Biostatistique*. Chicoutimi, Québec: Gaëtan Morin Éditeur. 850 p
- Schiffleithner V, Essl F (2016) Is it worth the effort? Spread and management success of invasive alien plant species in a Central European National Park. *NeoBiota* 31:43–61
- Stannard M, Crowder W (2003) Reed canarygrass. *Phalaris arundinacea* L. Pullman, WA: U.S. Department of Agriculture, Natural Resources Conservation Service. 4 p
- StataCorp (2013) *Stata* 13.1. College Station, TX: StataCorp
- Tabak NM, von Wettberg E (2008) Native and introduced jewelweeds of the Northeast. *Northeast Nat* 15:159–176
- Tanner RA, Jin L, Shaw R, Murphy ST, Gange AC (2014) An ecological comparison of *Impatiens glandulifera* Royle in the native and introduced range. *Plant Ecol* 215:833–843
- Tanner RA, Varia S, Eschen R, Wood S, Murphy ST, Gange AC (2013) Impacts of an invasive non-native annual weed, *Impatiens glandulifera*, on above- and below-ground invertebrate communities in the United Kingdom. *PLoS ONE* 8:e67271
- Thijs KW, Brys R, Verboven HAF, Hermy M (2012) The influence of an invasive plant species on the pollination success and reproductive output of three riparian plant species. *Biol Invasions* 14:355–365
- Vilà M, Bartomeus I, Dietzsch AC, Petanidou T, Steffan-Dewenter I, Stout JC, Tscheulin T (2009) Invasive plant integration into native plant–pollinator networks across Europe. *Proc R Soc Lond B Biol Sci* 276:3887–3893
- Wadsworth RA, Collingham YC, Willis SG, Huntley B, Hulme PE (2000) Simulating the spread and management of alien riparian weeds: are they out of control? *J Appl Ecol* 37(Suppl 1):28–38

Received February 15, 2017, and approved April 25, 2017.

Associate Editor for this paper: John A. Randall, University of California, Davis.